







Male Pattern Baldness Norwood Classifications					
2 Receding Hairline		3 Generalized frontal thinning		4 Frontal area & crown balding	
5 Top of Scalp & crown balding		6 Extensive hair loss with limited, yet viable donor area		7 Severe hair loss, only rim of hair remains	

Please indicate the(number of) persons with and without male pattern hair loss:	Please fill here the correct answers ↓ ↓	
	YES Male pattern Hair loss	NO Male pattern hair loss
FAMILY OF FATHER		
Grand-father (father of father) ?		
Uncles (brothers of father)?:		
Aunts (sisters of father)?		
FAMILY OF MOTHER		
Grand-father (father of mother)		
Uncles (brothers of mother)? :		
Aunts (sisters of mother)?		
YOUR FIRST FAMILY		
Father		
Mother		
Brothers?		
Sisters?		
YOUR SECOND FAMILY		
Yourself?	1	
Sons?		
Daughters ?		
TOTAL:		

⇒⇒

Treatment:

Racial predisposition to hair loss and baldness

Research has shown that Caucasian men are more likely to suffer from male pattern baldness - up to 80 percent of all white men will lose their hair in part by the time they are 60 years old. While Chinese men are the least likely to lose their hair - only 40 percent of all Chinese men will lose their hair by the age of 60.